

# Universality Class of $O(N)$ Models

**Adrian Patrascioiu**

Physics Department, University of Arizona  
Tucson, AZ 85721, U.S.A.

and

**Erhard Seiler**

Max-Planck-Institut für Physik  
– Werner-Heisenberg-Institut –  
Föhringer Ring 6, 80805 Munich, Germany

## **Abstract**

We point out that existing numerical data on the correlation length and magnetic susceptibility suggest that the two dimensional  $O(3)$  model with standard action has critical exponent  $\eta = 1/4$ , which is inconsistent with asymptotic freedom. This value of  $\eta$  is also different from the one of the Wess-Zumino-Novikov-Witten model that is supposed to correspond to the  $O(3)$  model at  $\theta = \pi$ .

Ever since the seminal paper of Kosterlitz and Thouless [1] it has been accepted that in two dimensions ( $2D$ ) there is a fundamental difference between the Abelian  $O(2)$  and the non-Abelian  $O(3)$  models: while the first one undergoes a transition (KT) from a phase with exponential decay at high temperature  $T$  ( $T = 1/\beta$ ) to a massless one at low temperature, the latter is in the high temperature phase for any  $\beta < \infty$ . While the first conjecture was proven rigorously by Fröhlich and Spencer [2], the second one remains unproven.

We have advanced several arguments [3, 4, 5] against this accepted belief and in our opinion *all*  $O(N)$  models undergo a KT-like transition at some finite  $\beta$ . Since however our arguments do not constitute a rigorous demonstration, it may be useful to get a glimpse at what the truth may be via existing numerical data. Our observation pertains to the value of the critical exponent  $\eta$ , defined as follows: let  $\xi$  be the correlation length and  $\chi$  the magnetic susceptibility. Then  $\eta$  can be defined by the following asymptotic statement:

$$\chi \propto \xi^{2-\eta} \quad (1)$$

i.e.  $\ln \chi - (2 - \eta) \ln \xi$  should go to a constant as  $\xi \rightarrow \infty$ .




Figure 1:  $O(2)$  model, test of  $\eta = 1/4$

For the  $O(2)$  model certain non-rigorous renormalization group arguments

advanced by Kosterlitz [6] predict that  $\eta = 1/4$ . In Tab.1 we collect the data from [7] and [8]. We could plot  $\ln \chi$  vs.  $\ln \xi$  and would see a nice straight line with a slope close to 1.75. In fact a fit of  $\ln \chi$  to a linear function of  $\ln \xi$  yields a slope of  $2 - \eta = 1.72$  with a good fit quality, if the errors on both  $\xi$  and  $\chi$  are taken into account. It is, however, much more significant to look at the deviations from the predicted slope by plotting  $\ln \chi - 1.75 \ln \xi$  vs.  $\ln \xi$ , together with the least square fit to a constant. This is done in Fig.1; the data of [7] are represented by crosses, the ones of [8] by squares. It can be seen that the data still show a tendency to decrease, corresponding to the fact that the asymptotic region has not yet been reached and the fits published in the literature always produced an  $\eta > 1/4$ , as did ours. But from Fig.1 one can also see a tendency of the data to flatten with increasing  $\xi$ , in accordance with the theoretical prediction of  $\eta = 1/4$ . The error bars shown in Fig.1 correspond to adding the errors on  $\ln \chi$  to 1.75 times the errors on  $\ln \xi$  (probably an overestimation as the errors on  $\chi$  and  $\xi$  are correlated).

According to the accepted scenario, the behavior of the  $O(3)$  model should be vastly different; indeed perturbative renormalization group arguments predict the following asymptotic behavior (“asymptotic scaling”):

$$\xi \propto \frac{\exp(2\pi\beta)}{\beta} \quad (2)$$

$$\chi \propto \frac{\exp(4\pi\beta)}{\beta^4} \quad (3)$$

From this it follows that for  $\xi \rightarrow \infty$  one should observe  $\ln \chi - 2 \ln \xi + 2 \ln \beta$  tending to a constant. To test this prediction, we use the published data of [9] as well as the unpublished ones due to [10], which are collected in Tab.2. In Fig.2 we plot  $\ln \chi - 2 \ln \xi + 2 \ln \beta$ , which should be constant according to the asymptotic scaling prediction. In Fig.3 we plot instead  $\ln \chi - 1.75 \ln \xi$  vs.  $\ln \xi$ , as we did for  $O(2)$ . Fig.3 also contains a least square fit to a constant. The data from [9] are represented by squares, the ones of [10] by crosses. The error bars are obtained, as in the case of  $O(3)$ , by adding the errors on  $\ln \chi$  and 1.75 times the errors on  $\ln \xi$ . Note that in both figures we are using the same scale as in Fig.1. Fitting  $\ln \chi$  to a linear function of  $\ln \xi$ , as we did for  $O(2)$ , gives a slope of  $2 - \eta = 1.74$ , again with a good fit quality ( $\chi^2 \approx 1$  per degree of freedom).

The following facts are visible from Figs 3 and 2:

- (1) The data are not consistent with asymptotic scaling inasmuch the points shown in Fig.2 are not consistent with a constant but show a strong decrease.
- (2) The data are consistent with a critical behavior with the same  $\eta = 1/4$  that was predicted by the KT theory for the  $O(2)$  model.

So we have to conclude that in fact the numerical data disagree with the predictions of asymptotic freedom and instead suggest that both  $O(2)$  and




Figure 2:  $O(3)$  model, test of asymptotic scaling

$O(3)$  have the same value of  $\eta$ . The data also indicate that the  $O(3)$  model with standard action  $\sum s(i) \cdot s(j)$  is not in the same universality class as the  $\theta = \pi$   $O(3)$  model, which is supposed to have the Wess-Zumino-Novikov-Witten model as its scaling limit [11]; conformal field theory arguments predict  $\eta=1$ , a prediction which a recent paper [12] claims to have verified numerically.

A note of caution: in principle  $\eta$  can also be determined by studying the finite size scaling of the susceptibility  $\chi(L)$ . Ideally this study should be performed at  $\beta_{crt}$ ; in practice though, it suffices to place oneself in a regime where  $\xi \gg L$ . One could ask then what value of  $\eta$  would come out of such a determination. The answer is that one must be careful and use a large enough  $L$ . Indeed, in this type of problems, besides the correlation length  $\xi$ , there is a second important length  $\xi_{PT}$ , the distance over which the system is well ordered. From perturbation theory (PT) one knows that for the  $O(3)$  model this second length is  $O(\exp(\pi\beta))$ . Consequently, if one uses periodic boundary conditions (b.c.) and an  $L$  and  $\beta$  such that  $\xi_{PT} \gg L$ , then the system will behave according to PT and one will find the perturbative value of  $\eta$  which is  $1/\pi\beta + O(\beta^{-2})$ . As we explained in a recent paper [14], a clear indication that such a determination cannot be trusted to reflect the true thermodynamic




Figure 3:  $O(3)$  model, test of  $\eta = 1/4$

value of  $\eta$  is the dependence of  $\eta$  upon the b.c. employed. Indeed, as we discussed in ref.[14], in the limit  $L \rightarrow \infty$ , the value of  $\eta$  must be the same if we use periodic b.c. and freeze two spins situated a distance  $L/2$  apart in any arbitrary positions; this then gives a criterion for deciding whether one has used a sufficiently large  $L$  and whether the determination of  $\eta$  is trustworthy.

Finally we should like to make a remark about the large  $N$ -limit of the  $O(N)$  model, which is often cited as evidence for the standard picture of  $\beta_{crt} = \infty$  for all  $N > 2$ . As we pointed out in various places (see in particular [15]), this reasoning is faulty. It is true that at fixed  $\tilde{\beta} \equiv \beta/N$  both  $\xi$  and  $\chi$  converge as  $N \rightarrow \infty$  to the values of the spherical model, hence in that limit  $\eta = 0$  up to log corrections. But, as we pointed out in [15], the approach to the limit is nonuniform, i.e. it becomes slower and slower as  $\tilde{\beta} \rightarrow \infty$  or  $\xi \rightarrow \infty$ . With increasing  $N$ ,  $\ln \chi$  as a function of  $\ln \xi$  will approach the spherical limit ( $N = \infty$ ) form, but at any fixed  $N$ , for large enough  $\beta$ , we expect that it will fall away from that form and that  $\ln \chi - 1.75 \ln \xi$  will approach a constant for  $\beta \rightarrow \infty$ .

We are indebted to S.Caracciolo, R.G.Edwards, A.Pelissetto and A.D.Sokal for allowing us to use their data prior to publication; these data form the basis

of the papers [13] and will appear in full in subsequent papers by [16].

## References

- [1] J.M.Kosterlitz and D.J.Thouless, *J. Phys. (Paris)* **32** (1975) 581.
- [2] J.Fröhlich and T.Spencer, *Commun.Math.Phys.* **81** (1981) 527.
- [3] A.Patrascioiu, *Existence of Algebraic Decay in non-Abelian Ferromagnets*, University of Arizona preprint AZPH-TH/91-49.
- [4] A.Patrascioiu and E.Seiler, *Percolation Theory and the Existence of a Soft Phase in 2D Spin Models*, *Nucl.Phys.B.(Proc. Suppl.)* **30** (1993) 184.
- [5] A.Patrasciou and E.Seiler, *Phys.Rev.Lett.* **74** (1995) 1920.
- [6] J.M.Kosterlitz, *J.Phys.* **C6** (1974) 1046.
- [7] U.Wolff *Nucl.Phys.* **B334** (1990) 581.
- [8] R.Gupta and C.F.Baillie, *Phys.Rev.B* **45** (1992) 2883.
- [9] J.Apostolakis, C.F. Baillie and G.F.Fox, *Phys.Rev.D* **43** (1990) 2687.
- [10] S.Caracciolo, R.G.Edwards, A.Pelissetto and A.D.Sokal, private communication.
- [11] I.Affleck *Phys.Rev.Lett.* **66** (1991) 2429.
- [12] W.Bietenholz, A.Pochinsky and U-J.Wiese, *Meron-Cluster Simulation of the  $\theta$ -Vacuum in 2-d  $O(3)$ -Model*, MIT preprint CPT 2433, hep-lat/9505019.
- [13] S.Caracciolo, R.G.Edwards, A.Pelissetto and A.Sokal, *Phys.Rev.Lett.* **74** (1995) 2969;  
– *Nucl.Phys.B (Proc.Suppl.)* **42** (1995) 752;  
– *Asymptotic Scaling in the Two-Dimensional  $O(3)$   $\sigma$ -Model at Correlation Length  $10^5$* , hep-lat/9411009, to appear in *Phys. Rev. Lett.*
- [14] A.Patrascioiu and E.Seiler, *Super-Instantons, Perfect Actions, Finite Size Scaling and the Continuum Limit*, preprint MPI-PhT/95-71, AZPH TH/95-17, hep-lat 9507018.
- [15] A.Patrascioiu and E.Seiler, *Nucl.Phys.* **B 443** (1995) 596.
- [16] S.Caracciolo, R.G.Edwards, A.Pelissetto and A.Sokal, in preparation.

**Tab.1a:**  $\chi$  and  $\xi$  for the  $O(2)$  model on thermodynamic lattices ( $L/\xi \geq 7$ ).  
Data are taken from [7].

| $\beta$ | $\xi$     | $\chi$     |
|---------|-----------|------------|
| .91     | 9.36(5)   | 134.17(45) |
| .92     | 10.69(8)  | 162.68(61) |
| .93     | 12.03(6)  | 201.29(53) |
| .94     | 13.50(9)  | 249.90(80) |
| .95     | 15.61(10) | 319.6(1.1) |
| .96     | 18.08(13) | 414.1(1.7) |
| .97     | 21.66(13) | 554.9(2.9) |
| .98     | 26.37(19) | 764.6(3.0) |
| .99     | 31.78(21) | 1092.(4.)  |
| 1.00    | 40.20(20) | 1604.(4.)  |
| 1.02    | 69.27(59) | 4170.(19.) |

**Tab.1b:**  $\chi$  and  $\xi$  for the  $O(2)$  model on thermodynamic lattices ( $L/\xi \geq 7$ ).  
Data are taken from [8].

| $\beta$  | $\xi$     | $\chi$     |
|----------|-----------|------------|
| .9090909 | 9.32(2)   | 132.41(55) |
| .9345794 | 12.61(2)  | 220.84(86) |
| .9478673 | 15.23(5)  | 302.19(89) |
| .9615385 | 18.70(20) | 433.7(2.2) |
| .9708738 | 21.80(20) | 564.1(2.7) |
| .9756098 | 23.90(20) | 665.0(4.0) |
| .9803922 | 26.20(20) | 779.0(5.0) |
| .9900990 | 31.70(30) | 1087.(10.) |
| 1.000000 | 40.40(40) | 1631.(12.) |
| 1.010101 | 51.30(90) | 2520.(24.) |
| 1.020408 | 70.(1.)   | 4258.(20.) |

**Tab.2a:**  $\chi$  and  $\xi$  for the  $O(3)$  model on thermodynamic lattices ( $L/\xi \geq 7$ ).  
Data are taken from [9].

| $\beta$ | $\xi$      | $\chi$      |
|---------|------------|-------------|
| 1.5     | 11.05(1)   | 176.4(2)    |
| 1.6     | 19.00(2)   | 448.4(7)    |
| 1.7     | 34.44(6)   | 1263.7(3.3) |
| 1.75    | 47.2(2)    | 2197.(15.)  |
| 1.8     | 64.5(5)    | 3823.(21.)  |
| 1.85    | 88.7(5)    | 6732.(25.)  |
| 1.9     | 122.7(1.1) | 11867.(62.) |

**Tab.2a:**  $\chi$  and  $\xi$  for the  $O(3)$  model on thermodynamic lattices ( $L/\xi \geq 7$ ).  
Data are taken from [10].

| $\beta$ | $\xi$     | $\chi$       |
|---------|-----------|--------------|
| 1.5     | 11.13(16) | 178.0(1.1)   |
| 1.55    | 14.69(15) | 281.5(1.7)   |
| 1.6     | 18.66(35) | 447.6(2.6)   |
| 1.65    | 25.21(13) | 742.8(2.0)   |
| 1.675   | 29.77(13) | 971.9(2.7)   |
| 1.7     | 34.66(13) | 1269.4(3.6)  |
| 1.725   | 40.21(25) | 1661.8(5.8)  |
| 1.75    | 47.20(19) | 2193.9(6.0)  |
| 1.775   | 55.23(25) | 2886.4(10.2) |
| 1.8     | 65.17(57) | 3839.7(23.8) |